

# A seven-bond coupling, ${}^7J(\text{PP})$ or through space coupling in the azine diphosphine $Z, Z\text{-PPh}_2\text{CH}_2\text{C}(\text{Bu}^t)=\text{N}-\text{N}=(\text{Bu}^t)\text{CCH}_2\text{PPh}_2$ : crystal structure of $Z, Z\text{-P}(\text{=O})\text{Ph}_2\text{CH}_2(\text{Bu}^t)=\text{N}-\text{N}=(\text{Bu}^t)\text{-CCH}_2\text{P}(\text{=O})\text{Ph}_2$

Sarath D. Perera, Bernard L. Shaw\*, Mark Thornton-Pett and Jonathan D. Vessey  
School of Chemistry, Leeds University, Leeds LS2 9JT (UK)

(Received December 24, 1992)

## Abstract

Analysis of the  ${}^{13}\text{C}\{^1\text{H}\}$  NMR spectrum of  $Z, Z\text{-PPh}_2\text{CH}_2\text{C}(\text{Bu}^t)=\text{N}-\text{N}=\text{C}(\text{Bu}^t)\text{CH}_2\text{PPh}_2$  (**1**) gives second order patterns for the  $\text{C}_{\text{ipso}}$ ,  $\text{C}_{\text{ortho}}$ ,  $\text{C}_{\text{meta}}$ ,  $\text{PCH}_2$  and  $\text{C}=\text{N}$  carbons caused by a non-zero seven-bond coupling,  ${}^7J(\text{PP})$ . Simulation of the spectra gives  ${}^7J(\text{PP}) = 4.8$  Hz. The corresponding diphosphine dioxide **2** gives first order  ${}^{13}\text{C}\{^1\text{H}\}$  spectra, i.e.  ${}^7J(\text{PP}) \sim 0$  Hz. Possible explanations for this unusually large value of  ${}^7J(\text{PP})$  are discussed. Crystals of **2** are orthorhombic, space group  $Pbca$ , with  $a = 1592.2(3)$ ,  $b = 1196.1(2)$ ,  $c = 1663.0(3)$  pm and  $Z = 4$ ; final  $R$  factor 0.0381 for 1780 observed reflections.

## Introduction


In a previous paper [1] we described the new azine diphosphine  $\text{PPh}_2\text{CH}_2\text{C}(\text{Bu}^t)=\text{N}-\text{N}=\text{C}(\text{Bu}^t)\text{CH}_2\text{PPh}_2$ , which was shown to have the  $Z, Z$  configuration by X-ray crystallography. When this azine diphosphine **1** was treated with  $[\text{M}(\text{CO})_4(\text{norbornadiene})]$  ( $\text{M} = \text{Cr}, \text{Mo}, \text{W}$ ) the norbornadiene was displaced to give  $[\text{M}(\text{CO})_4\{\text{PPh}_2\text{CH}_2\text{C}(\text{Bu}^t)=\text{N}-\text{N}=(\text{Bu}^t)\text{CCH}_2\text{PPh}_2\}]$  in which azine diphosphine is bonded in the  $E, Z$  configuration giving a nine-membered ring. These tetracarbonyl complexes were thermally unstable and when heated in benzene to  $c. 75^\circ\text{C}$ , the corresponding tricarbonyl complexes  $fac\text{-}[\text{M}(\text{CO})_3\{\text{PPh}_2\text{CH}_2\text{C}(\text{Bu}^t)=\text{N}-\text{N}=(\text{Bu}^t)\text{CCH}_2\text{PPh}_2\}]$  were formed with the  $E, Z$ -azine diphosphine acting as a tridentate ligand.

## Experimental

The general methods and instrumentation used were the same as those reported in recent papers from this laboratory [2]. The  ${}^{13}\text{C}\{^1\text{H}\}$  NMR spectra were recorded on a Bruker AM-400 spectrometer; for **1** the digitisation was increased to  $c. 0.05$  Hz/pt by recording the spectrum one resonance at a time. The spectra were simulated using standard Bruker software (PANIC programme).

\*Author to whom correspondence should be addressed.

The diphosphine **1** and the corresponding dioxide **2** were prepared as described previously [1].



## Single crystal X-ray diffraction analysis of **2**

All crystallographic measurements were carried out on a Stoe STADI4 diffractometer operating in the  $\omega/\theta$  scan mode using graphite monochromated  $\text{Mo K}\alpha$  radiation. The data-set was corrected for absorption semi-empirically using azimuthal psi scans.

The structure was determined by direct methods using SHELXS-86 [3] and was refined by full-matrix least-squares using the SHELX76 program system [4]. All non-hydrogen atoms were refined with anisotropic thermal parameters. The phenyl groups were treated as rigid bodies with idealised hexagonal symmetry ( $\text{C}-\text{C} = 139.5$  pm). All hydrogen atoms were included in calculated positions ( $\text{C}-\text{H} = 96$  pm) and were refined with an overall isotropic thermal parameter. The weight-

ing scheme  $w = [\sigma^2(F_o) + 0.0003(F_o)^2]^{-1}$  was used. For final non-hydrogen atomic coordinates see Table 3.

#### Crystal data

$C_{36}H_{42}N_2O_2P_2$ ,  $M = 596.69$ , orthorhombic, space group  $Pbca$ ,  $a = 1592.2(3)$ ,  $b = 1196.1(2)$ ,  $c = 1663.0(3)$  pm,  $U = 3.1671(11)$  nm<sup>3</sup>,  $Z = 4$ ,  $D_x = 1.25$  Mg m<sup>-3</sup>,  $\lambda(\text{Mo K}\alpha) = 71.069$  pm,  $\mu = 1.32$  cm<sup>-1</sup>,  $F(000) = 1272$ .

#### Data collection

$4.0 < 2\theta < 50.0^\circ$ , 3169 data collected, 1780 with  $I > 2.0$   $\sigma(I)$  considered observed,  $T = 80$  K.

#### Structure refinement

Number of parameters = 176,  $R = 0.0381$ ,  $R_w = 0.0390$ .

### Results and discussion

We have now studied the  $^{13}\text{C}\{^1\text{H}\}$  NMR spectrum of this unusual diphosphine in more detail than previously and find that some of the carbons of the phenyl rings, namely the *ipso*-, *ortho*- and *meta*-carbons, together with the resonances of the  $\text{PCH}_2$  and  $\text{C}=\text{N}$  carbons, give second order resonance patterns, which indicate a phosphorus–phosphorus coupling of a few hertz. The actual patterns are given in Fig. 1. We have calculated the patterns for these carbons using the measured values of  $J(\text{PC})$  and various values of  $J(\text{PP})$ . The best agreement between the observed and simulated (calculated) patterns is for  $J(\text{PP}) = 4.8$  Hz; as can be seen from Fig. 1, the agreement between the observed and simulated spectra is very good. The calculated spectra using other values of  $^7J(\text{PP})$ , e.g. 4.6 or 5.0 Hz, agree less with the observed spectra than does that using  $^7J(\text{PP}) = 4.8$  Hz.

In view of the unusually large long range coupling in **1** we have examined the  $^{13}\text{C}\{^1\text{H}\}$  NMR spectra of the corresponding diphosphine dioxide **2** (see Table 1). None of the  $^{13}\text{C}$  resonances showed second order features, indicating that  $^7J(\text{PP})$  is approximately zero. We have also determined the crystal structure of this diphosphine dioxide **2** by X-ray diffraction and established that it also has the *Z,Z* configuration (Fig. 2, see below).

The value of 4.8 Hz for  $^7J(\text{PP})$  is ostensibly that of a seven-bond coupling constant. In the extensive studies of Colquhoun and McFarlane [5] on diphosphines of type  $\text{Ph}_2\text{PXPPH}_2$  with X = a variety of bridging groups including  $-(\text{CH}_2)_x-$  ( $x = 1-3$ ), *cis* and *trans*  $-\text{CH}=\text{CH}-$ ,  $-\text{C}\equiv\text{C}-$  etc., the values of  $J(\text{PP})$  were measured by triple resonance experiments, e.g. for the ditertiary phosphines of type  $\text{Ph}_2\text{P}(\text{CH}_2)_x\text{PPh}_2$   $^2J(\text{PP}) = +125$  Hz ( $x = 1$ ),  $^3J(\text{PP}) = +33.6$  Hz ( $x = 2$ ) and  $^4J(\text{PP}) = 1.0$  Hz ( $x = 3$ ). Other workers have observed zero coupling constants for  $^4J(\text{PP})$  [6] and  $^6J(\text{PP})$  [7].




Fig. 1. Observed (left) and simulated (right) portions of the  $^{13}\text{C}\{^1\text{H}\}$  NMR spectrum of **1** in  $\text{CDCl}_3$  at 100.6 MHz. (a)  $\text{C}_{ipso}$ ; (b)  $\text{C}_{ortho}$ ; (c)  $\text{C}_{meta}$ ; (d)  $\text{C}=\text{N}$ ; (e)  $\text{PCH}_2$ .

Appreciable coupling constants  $^7J(\text{HH})$  and  $^7J(\text{FF})$  have been observed in fully conjugated systems, e.g. styrene [8] and fluorostyrenes [9]. A value of  $^7J(\text{PP})$  of 0.3 Hz has been reported for

TABLE 1.  $^{13}\text{C}\{^1\text{H}\}$  NMR data<sup>a</sup>

Compound	$\delta_c$ (ppm)	$\Delta\delta$ (ppb)	$^nJ$ (Hz)	$^{n'}J$ (Hz)	Assignment
1	168.1	2.4	5.0 ( $n=2$ )	1.6 ( $n'=5$ )	C=N
	139.8	22.9	18.2 ( $n=1$ )	-0.2 ( $n'=8$ )	$C_{ipso}$
	133.0	4.3	20.4 ( $n=2$ )	0.3 ( $n'=9$ )	$C_{ortho}$
	128.5				$C_{para}$
	128.3	1.9	6.7 ( $n=3$ )	0.0 ( $n'=10$ )	$C_{meta}$
	38.4				$\text{CMe}_3$
	28.8	40.8	22.5 ( $n=1$ )	-2.6 ( $n'=6$ )	$\text{CH}_2$
	28.3				$\text{CMe}_3$
	2	167.4		8.7 ( $n=2$ )	2.3 ( $n'=5$ )
134.8			100.1 ( $n=1$ )		$C_{ipso}$
131.2			3.0 ( $n=4$ )		$C_{para}$
130.5			9.5 ( $n=2$ )		$C_{ortho}$
128.2			11.9 ( $n=3$ )		$C_{meta}$
38.4					$\text{CMe}_3$
32.0			62.9 ( $n=1$ )		$\text{CH}_2$
28.3					$\text{CMe}_3$

<sup>a</sup>Recorded in 100.6 MHz in  $\text{CDCl}_3$ ;  $\Delta\delta = |\delta(^{13}\text{C}_P - \delta(^{12}\text{C}_P))|$ ;  $n$  is the order of the coupling to the proximate phosphorus and  $n'$  the order of the coupling to the more remote phosphorus.




Fig. 2. ORTEP [13] representation of the crystal structure of 2.

$(^i\text{Pr})_2\text{PCH}_2\text{P}(^i\text{Pr})(\text{CH}_2\text{CH}_2\text{P}(^i\text{Pr})\text{CH}_2\text{P}(^i\text{Pr})_2$  [10] but we are unable to find in the literature other values of  $^7J(\text{PP})$ .

The relatively large value for the  $^7J(\text{PP})$  in 1 may well be due to conformational control of the  $\text{PPh}_2$  groups by the tertiary butyl groups: the crystal structure of 1 shows the phosphine phenyl groups to be almost eclipsed with respect to the  $\text{CH}_2$  hydrogens (dihedral angle  $\text{C}(15)\text{--P}(8)\text{--C}(7)\text{--H}$  is  $14.7(3)^\circ$ ), an orientation also observed in the structure of 2. Long-range coupling

TABLE 2. Bond lengths (pm) and angles ( $^\circ$ ) for compound 2 with e.s.d.s in parentheses

O(1)–P(1)	148.5(3)	C(3)–P(1)	182.6(5)
C(4)–P(1)	180.8(3)	C(10)–P(1)	181.3(3)
C(1)–N(1)	128.4(4)	N(1)–N(1*)	142.8(5)
C(2)–C(1)	153.1(6)	C(3)–C(1)	151.5(5)
C(21)–C(2)	153.0(6)	C(22)–C(2)	152.5(6)
C(23)–C(2)	153.4(6)		
C(3)–P(1)–O(1)	115.0(2)	C(4)–P(1)–O(1)	112.5(2)
C(4)–P(1)–C(3)	107.3(2)	C(10)–P(1)–O(1)	112.7(2)
C(10)–P(1)–C(3)	105.0(2)	C(10)–P(1)–C(4)	103.6(2)
C(1)–N(1)–N(1*)	112.8(3)		
C(2)–C(1)–N(1)	117.6(3)	C(3)–C(1)–N(1)	124.1(3)
C(3)–C(1)–C(2)	118.2(3)		
C(21)–C(2)–C(1)	111.8(3)	C(22)–C(2)–C(1)	110.8(3)
C(22)–C(2)–C(21)	108.7(4)	C(23)–C(2)–C(1)	106.8(3)
C(23)–C(2)–C(21)	109.1(3)	C(23)–C(2)–C(22)	109.6(3)
C(1)–C(3)–P(1)	112.2(3)	C(5)–C(4)–P(1)	121.6(2)
C(9)–C(4)–P(1)	118.1(2)	C(11)–C(10)–P(1)	117.9(2)
C(15)–C(10)–P(1)	121.9(2)		

N(1\*) is related to N(1) by the symmetry operator ( $-x, -y, 1.0-z$ ).

to  $\text{P}^{\text{III}}$  nuclei (but not  $\text{P}^{\text{V}}$  nuclei) is known to be particularly dependent upon the dihedral angles between bonds and the  $\text{P}^{\text{III}}$  lone pair [11], and it may be that the conformation of the  $\text{PPh}_2$  groups imposed by the tertiary butyl groups fortuitously aligns the  $\text{P}^{\text{III}}$  lone pairs for good transmission of the coupling through the backbone. An alternative explanation involves a direct interaction between the phosphorus and the  $\pi$ -system of the azine backbone: since the coupling is only observed for the diphosphine 1 and not for the corresponding dioxide 2, this implies that the interaction involves the phosphorus lone pair. Furthermore, it is

TABLE 3. Atom coordinates ( $\times 10^4$ ) for compound **2** with e.s.d.s in parentheses

	x	y	z
P(1)	720.6(4)	1130.1(6)	6763.6(4)
O(1)	670(1)	2233(2)	6355(1)
N(1)	-382(1)	311(2)	5028(1)
C(1)	-595(2)	520(2)	5757(2)
C(2)	-1424(2)	1141(3)	5895(2)
C(21)	-1369(2)	1950(3)	6606(2)
C(22)	-1676(2)	1797(3)	5147(2)
C(23)	-2091(2)	249(3)	6077(2)
C(3)	-103(2)	139(2)	6487(2)
C(4)	1717(1)	443(1)	6599(1)
C(5)	1886(1)	-614(1)	6915(1)
C(6)	2688(1)	-1072(1)	6843(1)
C(7)	3321(1)	-474(1)	6454(1)
C(8)	3152(1)	583(1)	6138(1)
C(9)	2350(1)	1041(1)	6211(1)
C(10)	656(1)	1244(1)	7849(1)
C(11)	781(1)	2294(1)	8194(1)
C(12)	800(1)	2408(1)	9028(1)
C(13)	695(1)	1472(1)	9518(1)
C(14)	570(1)	422(1)	9174(1)
C(15)	550(1)	308(1)	8339(1)

also possible that in some way a direct phosphorus-phosphorus lone pair interaction is contributing towards coupling. It has been shown that the radical cation  $\text{Ph}_2\text{PCH}_2\text{CH}_2\text{PPh}_2^+$  has two equivalent phosphorus nuclei and that the diphosphine is in the *cisoid* fashion with the lone pair and singly occupied orbitals on the phosphorus interacting [12]. As mentioned above,  $^3J(\text{PP})$  in  $\text{Ph}_2\text{PCH}_2\text{CH}_2\text{PPh}_2$  is 33.6 Hz, but in *cis*- $\text{Ph}_2\text{PCH}=\text{CHPPh}_2$   $^3J(\text{PP})$  is 105.5 Hz whereas in *trans*- $\text{Ph}_2\text{PCH}=\text{CHPPh}_2$  it is only 13.4 Hz. It is therefore possible that the lone pairs on phosphorus are interacting directly in *cis*- $\text{PPh}_2\text{CH}=\text{CHPPh}_2$  to give the large value of  $^3J(\text{PP})$ .

*Crystal structure of Z,Z-Ph<sub>2</sub>P(=O)CH<sub>2</sub>C(Bu')=N-N=C(Bu')CH<sub>2</sub>P(=O)Ph<sub>2</sub> (2)*

The crystal structure of the azine diphosphine dioxide **2** is shown in Fig. 2, with selected bond lengths and

angles in Table 2 and atom coordinates in Table 3. The crystal structure of **2** shows that the arrangement around both C=N bonds is *Z*. The bond lengths of the backbone are very similar to those reported for the corresponding diphosphine **1** [1] except that the bond lengths between phosphorus and carbon are slightly less in **2** than in **1**, as would be expected.

### Acknowledgements

We thank the SERC for postdoctoral research fellowships (to S.D.P. and J.D.V.) and for other support.

### References

- 1 S. D. Perera, B. L. Shaw and M. Thornton-Pett, *J. Chem. Soc., Dalton Trans.*, (1992) 1469.
- 2 B. L. Shaw and J. D. Vessey, *J. Chem. Soc., Dalton Trans.*, (1991) 3303.
- 3 G. M. Sheldrick, *SHELXS 86*, program system for X-ray structure solution, University of Göttingen, Germany, 1976.
- 4 G. M. Sheldrick, *SHELX 76*, program system for X-ray structure determination, University of Cambridge, UK, 1976.
- 5 I. J. Colquhoun and W. McFarlane, *J. Chem. Soc., Dalton Trans.*, (1982) 1915.
- 6 S. O. Grim, R. C. Barth, J. D. Mitchell and J. D. Gaudio, *Inorg. Chem.*, 16 (1977) 1776.
- 7 R. B. King and J. C. Cloyd, *Inorg. Chem.*, 14 (1975) 1550.
- 8 M. Barfield, C. J. MacDonald, R. I. Peat and W. F. Reynolds, *J. Am. Chem. Soc.*, 93 (1971) 4195.
- 9 D. A. Burgess, I. D. Rae and J. D. Snell, *Aust. J. Chem.*, 30 (1977) 543.
- 10 H. Sommer, S. Hietkamp and O. Stelzer, *Chem. Ber.*, 117 (1984) 3414.
- 11 L. D. Quin, M. J. Gallagher, G. T. Cunkle and D. B. Chesnut, *J. Am. Chem. Soc.*, 102 (1980) 3136.
- 12 C. J. Rhodes and M. C. R. Symons, *J. Chem. Soc., Chem. Commun.*, (1989) 1393.
- 13 C. K. Johnson, *ORTEP II*, Rep. ORNL-5138, Oak Ridge National Laboratory, TN, USA, 1976.